

Plant microRNA Purification Kit

Product # 54700

Product Insert

Introduction

Norgen's Plant microRNA Purification Kit provides a rapid method for the isolation and purification of small RNA molecules (< 200 nt) from cultured plant cells or plant tissues. These small RNAs include regulatory RNA molecules such as microRNA (miRNA) and short interfering RNA (siRNA), as well as tRNA and 5S rRNA. Small RNA molecules are often studied due to their ability to regulate gene expression. miRNAs and siRNAs are typically 20-25 nucleotides long, and regulate gene expression by binding to mRNA molecules and affecting their stability or translation. The small RNA molecules isolated using Norgen's Plant microRNA Purification Kit can be used in various downstream applications relating to gene regulation and functional analysis, including RT-PCR, miRNA sequencing, northern blotting and microarray analysis.

Norgen's Purification Technology

Purification is based on spin column chromatography using Norgen's proprietary resin as the separation matrix. Norgen's resin binds RNA in a manner that depends on ionic concentrations. The small RNA molecules are preferentially purified from other cellular components such as ribosomal RNA without the use of phenol or chloroform. The process involves the use of two different spin columns: the **Large RNA Removal Column** and the **microRNA Enrichment Column** (please see flow chart on page 3). Briefly, the cells or tissues of interest are lysed using the provided Lysis Buffer C, the lysate is spun through the provided the Filter Column, and then ethanol is added to the clarified lysate. The lysate is then applied to the Large RNA Removal Column, and the larger RNA molecules will bind to the resin in the spin column while the smaller RNA species will pass through into the flowthrough. Ethanol is then added to the flowthrough, and the sample is applied to the microRNA Enrichment Column. The small RNA molecules will then bind to the resin, and any impurities are removed through a series of washes with the provided Wash Solution A. The small RNA molecules are then eluted using the Elution Solution, and are ready for use in various applications.

Specifications

Kit Specifications	
Maximum Column Binding Capacity	50 µg
Maximum Column Loading Volume	650 µL
Minimum Elution Volume	20 µL
Size of RNA Purified	< 200 nt
Amount of Starting Material: Plant Tissues	100 mg 5 x 10 ⁶ cells
Time to Complete 10 Purifications	25 minutes

Advantages

- Fast and easy processing using a rapid spin-column format
- No phenol or chloroform extractions
- Isolate all small RNA molecules (<200 nt)
- Minimal contamination from large RNA molecules and genomic DNA
- High quality small RNA can be used in various downstream applications

Kit Components

Component	Product # 54700 (25 preps)
Lysis Buffer C	30 mL
Wash Solution A	18 mL
Elution Solution A	6 mL
Filter Columns	25
Large RNA Removal Columns	25
microRNA Enrichment Columns	25
Collection Tubes	50
Elution tubes (1.7 mL)	50
Product Insert	1

Storage Conditions and Product Stability

All solutions should be kept tightly sealed and stored at room temperature. These reagents should remain stable for at least 1 year in their unopened containers.

Precautions and Disclaimers

This kit is designed for research purposes only. It is not intended for human or diagnostic use. Ensure that a suitable lab coat, disposable gloves and protective goggles are worn when working with chemicals. For more information, please consult the appropriate Material Safety Data Sheets (MSDSs). These are available as convenient PDF files online at www.norgenbiotek.com.

The **Lysis Buffer C** contains guanidinium salts, and should be handled with care. Guanidinium salts form highly reactive compounds when combined with bleach, thus care must be taken to properly dispose of any of these solutions

Customer-Supplied Reagents and Equipment

- Benchtop microcentrifuge
- 96 - 100% ethanol
- RNase-free microcentrifuge tubes
- Liquid nitrogen
- Mortar and pestle

Working with RNA


RNases are very stable and robust enzymes that degrade RNA. Autoclaving solutions and glassware is not always sufficient to actively remove these enzymes. The first step when preparing to work with RNA is to create an RNase-free environment. The following precautions are recommended as your best defense against these enzymes.

- The RNA area should be located away from microbiological work stations
- Clean, disposable gloves should be worn at all times when handling reagents, samples, pipettes, disposable tubes, etc. It is recommended that gloves are changed frequently to avoid contamination
- There should be designated solutions, tips, tubes, lab coats, pipettes, etc. for RNA only
- All RNA solutions should be prepared using at least 0.05% DEPC-treated autoclaved water or molecular biology grade nuclease-free water
- Clean all surfaces with commercially available RNase decontamination solutions
- When working with purified RNA samples, ensure that they remain on ice during downstream applications


Flow Chart

Procedure for Purifying Small RNA Molecules using Norgen's Plant microRNA Purification Kit



Lyse cells or tissue using **Lysis Buffer C**




SPIN




SPIN



SPIN



SPIN



Purified Small RNA (miRNA, siRNA, tRNA, 5S rRNA, etc)

Procedures

All centrifugation steps are carried out in a benchtop microcentrifuge. Various speeds are required for different steps, so please check your microcentrifuge specifications to ensure that it is capable of the proper speeds. All centrifugation steps are performed at room temperature. The correct rpm can be calculated using the formula:

$$RPM = \sqrt{\frac{RCF}{(1.118 \times 10^{-5}) (r)}}$$

where *RCF* = required gravitational acceleration (relative centrifugal force in units of g); *r* = radius of the rotor in cm; and *RPM* = the number of revolutions per minute required to achieve the necessary *g*-force.

Section 1. Preparation of Lysate from Various Cell Types

All centrifugation steps are carried out in a benchtop microcentrifuge at 14,000 x g (~14,000 RPM) except where noted. All centrifugation steps are performed at room temperature.

Notes Prior to Use

- A variable speed centrifuge should be used for maximum kit performance. If a variable speed centrifuge is not available a fixed speed centrifuge can be used, however reduced yields may be observed.
- Ensure that all solutions are at room temperature prior to use.
- There are 2 different spin columns provided with this kit; the **Large RNA Removal Column** and the **microRNA Enrichment Column**. Ensure that the correct column is used for each step of the procedure.
- Prepare a working concentration of the **Wash Solution A** by adding 42 mL of 96 – 100 % ethanol (provided by the user) to the supplied bottle containing the concentrated **Wash Solution**. This will give a final volume of 60 mL. The label on the bottle has a box that may be checked to indicate that the ethanol has been added.
- The recommended input is 100 mg of plant tissue or 5×10^7 cells. However it can be reduced (to 30 mg) or increased (up to 150 mg) depending on customer's optimization based on a sample type.
- Both fresh and frozen plant tissues can be used for this protocol. Samples should be flash-frozen in liquid nitrogen and transferred immediately to a -70°C freezer for long-term storage. Do not allow frozen tissues to thaw prior to grinding with the mortar and pestle in order to ensure that the integrity of the RNA is not compromised.
- It is important to work quickly during this procedure.

1. Cell Lysate Preparation

- a. Transfer ≤ 100 mg of plant tissue or 5×10^7 plant cells (see Notes Prior to Use) into a mortar that contains an appropriate amount of liquid nitrogen to cover the sample. Grind the sample into a fine powder using a pestle in liquid nitrogen.

Note: If stored frozen samples are used, do not allow the samples to thaw before transferring to the liquid nitrogen.

- b. Allow the liquid nitrogen to evaporate, without allowing the tissue to thaw.
- c. Transfer the powder to a 1.5 mL centrifuge tube (not provided)
- d. Add 600 μ L of **Lysis Buffer C** and vortex vigorously for 30 seconds.
- e. Assemble a Filter Column (**clear O-ring**) with one of the provided collection tubes.
- f. Pipette the lysate into the Filter Column and spin for 2 minutes at 14,000 x g (~14,000 RPM).

- g. Transfer only the clear supernatant from the flow-through into a RNAase-free microcentrifuge tube (not provided) using a pipette. Note the volume of the supernatant/lysate.

Note: Ensure that only the clear supernatant is transferred, avoiding any of the debris at the bottom of the collection tube.

- h. Add a volume of 96-100% ethanol (provided by the user) that is a half of the lysate volume (50 μ L of ethanol is added to every 100 μ L of lysate). Vortex to mix. **Proceed to Step 2.**

2. Large RNA Removal

- a. Retrieve a **Large RNA Removal Column** assembled with a collection tube.
- b. Apply the lysate with the ethanol (from step 1) onto the column and centrifuge for 1 minute. **Retain the flowthrough, which contains the small RNA species.** If small RNA-depleted RNA is to be isolated, retain the column and proceed to the Optional Large RNA Purification Protocol (Appendix A). Otherwise, discard the column.

Note: The flowthrough contains the small RNA, thus ensure that this fraction is not discarded.

3. Small RNA Capture

- a. Add 1 volume of 96 – 100% ethanol (provided by the user) to the flowthrough collected in **Step 2b** (100 μ L of ethanol is added to every 100 μ L of flowthrough). Mix by vortexing for 10 seconds.
- b. Assemble a **microRNA Enrichment Column** with one of the provided collection tubes.
- c. Apply half of the lysate mix with ethanol onto the column and centrifuge for 1 minute.
- d. Discard the flowthrough and reassemble the spin column with the collection tube.
- e. Repeat steps **3c** and **3d** to complete the capture of the small RNA.

4. Column Wash

- a. Apply 400 μ L of **Wash Solution A** to the **microRNA Enrichment Column** and centrifuge for 1 minute.

Note: Ensure the entire wash solution has passed through into the collection tube by inspecting the column. If the entire wash volume has not passed, spin for an additional minute.

- b. Discard the flowthrough and reassemble the spin column with the collection tube.
- c. Repeat steps **4a** and **4b** to wash column a second time.
- d. Wash column a third time by adding another 400 μ L of **Wash Solution A** and centrifuging for 1 minute.
- e. Discard the flowthrough and reassemble the spin column with its collection tube.
- f. Spin the column for 2 minutes in order to thoroughly dry the resin. Discard the collection tube.

5. Small RNA Elution

- a. Place the **microRNA Enrichment Column** into a fresh 1.7 mL Elution tube provided with the kit.
- b. Add 20 μ L of **Elution Solution A** to the column.

- c. Centrifuge for 2 minutes at **200 x g (~2,000 RPM)**, followed by 1 minute at **14,000 x g (~14,000 RPM)**. Note the volume eluted from the column. If the entire volume has not been eluted, spin the column at 14,000 x g (~14,000 RPM) for 1 additional minute.

Note: For maximum RNA recovery, it is recommended that a second elution be performed into a separate microcentrifuge tube (Repeat **Steps 5b and 5c**).

6. Storage of RNA

The purified RNA sample may be stored at -20°C for a few days. It is recommended that samples be placed at -70°C for long term storage.

Appendix A: Optional Large RNA Purification Protocol

1. Column Wash

- a. Reassemble the **Large RNA Removal Column** with the collection tube used in Step **2b**
- b. Apply 400 μL of **Wash Solution A** to the **Large RNA Removal Column** and centrifuge for 1 minute.

Note: Ensure that the entire Wash Solution has passed through into the collection tube by inspecting the column. If the entire wash volume has not passed, spin for an additional minute.

- c. Discard the flowthrough and reassemble the spin column with the collection tube.
- d. Repeat steps **1b** and **1c** to wash the column a second time.
- e. Wash column a third time by adding another 400 μL of **Wash Solution A** and centrifuging for 1 minute.
- f. Discard the flowthrough and reassemble the spin column with its collection tube.
- g. Spin the column for 2 minutes in order to thoroughly dry the resin. Discard the collection tube.

2. Large RNA Elution

- a. Place the **Large RNA Removal Column** into a fresh 1.7 mL Elution tube provided with the kit.
- b. Add 50 μL of **Elution Solution A** to the column.
- c. Centrifuge for 2 minutes at **200 x g (~2,000 RPM)**, followed by 1 minute at **14,000 x g (~14,000 RPM)**. Note the volume eluted from the column. If the entire volume has not been eluted, spin the column at 14,000 x g (~14,000 RPM) for 1 additional minute.

Note: For maximum RNA recovery, it is recommended that a second elution be performed into a separate microcentrifuge tube (Repeat **Steps 2b and 2c**).

3. Storage of RNA

The purified RNA sample may be stored at -20°C for a few days. It is recommended that samples be placed at -70°C for long term storage.

Troubleshooting Guide

Problem	Possible Cause	Solution and Explanation
Poor RNA Recovery	Incomplete lysis of cells or tissue	Ensure that the appropriate homogenization step and amount of Lysis Buffer C was used for the amount of cells or tissue. Incubate the sample with the Lysis Buffer C at 55°C for 5 minutes.
	Large RNA Removal Column has become clogged	Do not exceed the recommended amounts of starting materials. The amount of starting material may need to be decreased if the column shows clogging below the recommended levels. See also “Clogged Column” below.
	An alternative elution solution was used	It is recommended that the Elution Solution A supplied with this kit be used for maximum RNA recovery.
	Low RNA content	Different tissues and cells have different RNA contents. Some tissues may not contain small RNA at detectable levels when processing the small sample sizes required for this procedure.
	Flowthrough from the first binding step was discarded	The flowthrough from the binding step with the Large RNA Removal Column contains the small RNA molecules, thus ensure that it is not inadvertently discarded.
	Ethanol was not added to the flowthrough before binding to the microRNA Enrichment Column	Ensure that the appropriate amount of ethanol was added to the flowthrough from the first binding step before it is applied to the microRNA Enrichment Column. This is imperative in order to capture the small RNA molecules.
	Ethanol was not added to the Wash Solution	Ensure that 42 mL of 96 – 100 % ethanol is added to the supplied Wash Solution A prior to use.

Problem	Possible Cause	Solution and Explanation
Clogged Column	Insufficient solubilization of cells or tissues	Ensure that the appropriate amount of Lysis Buffer C was used for the amount of cells or tissue.
	Maximum number of cells or amount of tissue exceeds kit specifications	Refer to specifications to determine if amount of starting material falls within kit specifications.
	Centrifuge temperature too low	Ensure that the centrifuge remains at room temperature throughout the procedure. Temperatures below 15°C may cause precipitates to form that can cause the columns to clog.
RNA is Degraded	RNase contamination	RNases may be introduced during the use of the kit. Ensure proper procedures are followed when working with RNA. Please refer to “ <i>Working with RNA</i> ” at the beginning of this user guide.
	Procedure not performed quickly enough	In order to maintain the integrity of the RNA, it is important that the procedure be performed quickly.
	Improper storage of the purified RNA	For short term storage RNA samples may be stored at –20°C for a few days. It is recommended that samples be stored at –70°C for longer term storage.
	Frozen tissues or pellets were allowed to thaw prior to disruption	Tissue samples should be flash-frozen in liquid nitrogen and transferred immediately to a -70°C freezer for long-term storage. Do not allow frozen tissues to thaw prior to grinding with the mortar and pestle in order to ensure that the integrity of the RNA is not compromised.
RNA does not perform well in downstream applications	RNA was not washed three times with the provided Wash Solution	Traces of salt from the binding step may remain in the sample if the microRNA Enrichment Column is not washed three times with Wash Solution A. Salt may interfere with downstream applications, and thus must be washed from the column.
	Ethanol carryover	Ensure that the dry spin under the Column Wash procedure is performed, in order to remove traces of ethanol prior to elution. Ethanol is known to interfere with many downstream applications.

Problem	Possible Cause	Solution and Explanation
Genomic DNA contamination	Large amount of starting material used	Perform RNase-free DNaseI digestion on the RNA sample after elution to remove genomic DNA contamination.
Large RNA species present in elution	Improper amount of ethanol added to the lysate before binding to the Large RNA Removal Column	Ensure that the appropriate amount of ethanol was added to the lysate before it is applied to the Large RNA Removal Column. This is imperative in order to capture the large RNA molecules onto the column.
	Large amount of starting material used	Repeat purification using less starting material. Alternatively, the isolation procedure can be repeated using the elution as the input. The elution volume should first be adjusted to 300 μ L using the provided Lysis Solution. The procedure can then be followed as written in the manual, starting with the addition of ethanol, centrifuging the lysate in order to pellet any debris, and applying the clarified lysate to the Large RNA Removal Column. Repeating the procedure should result in the removal of the large, contaminating RNA species.

Related Products	Product #
100b RNA Ladder	15002
1kb RNA Ladder	15003
Plant/Fungi Total RNA Purification kit	25800, 31350, 31900
Plant RNA/DNA Purification Kit	24400

Technical Support

Contact our Technical Support Team between the hours of 8:30 and 5:30 (Eastern Standard Time) at (905) 227-8848 or Toll Free at 1-866-667-4362.

Technical support can also be obtained from our website (www.norgenbiotek.com) or through email at techsupport@norgenbiotek.com.

3430 Schmon Parkway, Thorold, ON Canada L2V 4Y6
 Phone: (905) 227-8848
 Fax: (905) 227-1061
 Toll Free in North America: 1-866-667-4362