


Hydroxystilbamidine

Product Information

Name :	Hydroxystilbamidine isethionate also known as FluoroGold®
Catalog Number :	UP544201 , 10mg UP544200, 50mg
Structure :	
Molecular Weight :	MW= 532.6
Soluble:	H2O or DMSO
Absorption / Emission :	$\lambda_{exc} \lambda_{em}$ (pH4.5) = 323 / 408 nm; EC: 33 000 M ⁻¹ cm ⁻¹ $\lambda_{exc} \lambda_{em}$ (membrane bound) = 350-395 / 530-600 nm $\lambda_{exc} \lambda_{em}$ (DNA) = 360 / ~450, ~625; EC: 27 000 M ⁻¹ cm ⁻¹



Storage: +4°C (-20 °C for long term) and protect from light, desiccated.

Hydroxystilbamidine is also in solution at 4% (FP-JW7290)

Introduction

Fluoro-Gold (Hydroxystilbamidine) has been used extensively as a retrograde tracer for neurons and also a histochemical stain, and is more flexible in terms of post-injection survival times, concentration range, tissue treatment and compatibility with other histochemical techniques.

Advantages of Fluoro-Gold when used alone...

- Intensively vibrant retrograde fluorescent tracer that is extremely sensitive and reliable
- Not taken up by undamaged fibers of passage
- Does not diffuse out of retrogradely labeled neurons
- Wide range of survival times from one day to over a month
- Stable fluorochrome that takes superlative colour and black and white photos
- Easy and safe – long shelf life- pure reliable source
- Can be pressure injected or iontophoresed
- Compatible with most other frequently used neuroanatomical techniques *

*such as immunofluorescence, immunocytochemistry, autoradiography, HRP histochemistry, paraffin embedded and other retrograde fluorescent tracers

Advantages of Fluoro-Gold when enhanced by the antibody.

- Substantially enhances visualization of Fluo-Gold
- Allows much more precise anatomical mapping
- Shows extremely fine anatomical structure including dendrites axons and terminals
- Permanent staining when used with enzyme amplification kits

Directions for use

Dry Fluoro-Gold should be kept in the dark at -4°C, or -20°C for long term storage, but is also stable at room temperature for well over six months. The dye in solution should be kept in the dark at 4 degrees celcius and will remain stable for at least six months.

Contact your local distributor
Info@fluoprobes.com
 Technical-support@fluoprobes.com
Order-online@fluoprobes.com

Fluoprobes, powered by



213 Avenue J.F. Kennedy - BP 1140
 93163 Montfermeil Cedex - France
 Tél. 04 70 03 88 55 - Fax 04 70 03 82 80

FT-T8135

Fluoro-Gold can be dissolved in distilled water or 0.9% saline, or utilized as a suspension in 0.2M neutral phosphate buffer.

Fluoro-Gold has been successfully used at concentrations ranging from 1-10%. Initially, a 4% concentration is advised. If undesirable necrosis occurs at the injection site, or labeling is too intense, reduce the concentration to a 2% solution.

A. Pressure Injection: this is probably the most frequently used mode of application. Volumes injected range from .05-1 ul, typically .1-2 ul.

B. Iontophoresis: discrete, small injection sites result from 4-10 second pulsed iontophoretic (+5 to +10ua/10min) application.

C. Crystal: a crystal of the tracer can be administered from the tip of a micro-pipette.

Good retrograde labeling has been observed with periods ranging from two days to two months. Survival periods of three to five days are typical. Long survival periods enhance filling of distal processes without diffusion of the dye from the cell.

Most any fixative, or no fixative, can be used, Phosphate neutral buffered saline containing 4% formaldehyde is frequently employed. Fixatives containing high concentrations of heavy metals (e.g. osmium, mercury) will quench the fluorescence, while high concentrations (over 1%) of glutaraldehyde may increase background fluorescence

Tissue containing Fluoro-Gold may be processed according to virtually any common histological technique. This includes cryostat sections of unfixed tissue (10 μ m), frozen sections of fixed tissue (20 μ m), and thin sections cut from tissue imbedded in either plastic (2-4 μ m) or paraffin (3-10 μ m). Frozen sections of fixed tissue are most frequently used.

At this point of processing, sections may be further processed for a second marker such as autoradiography, HRP histochemistry, immunocytochemistry, a second fluorescent tracer, fluorescent counterstain, etc.

Sections are typically mounted on gelatin-coated slides, air-dried, immersed in xylene, and coverslipped with non-fluorescent DPX plastic mounting media. Sections may be dehydrated with graded alcohols, unless this is not compatible with a second tracer. If Fluoro-Gold is to be combined with fluorescence immunocytochemistry, then sections are air-dried and directly coverslipped with neutral buffered glycerine (1:2).

Fluoro-Gold can be visualized with a fluorescence microscope using a wide band ultraviolet excitation filter (Excitation - 323 nm, Emission - 620 nm at neutral pH). A gold color is emitted when tissue has been processed with neutral pH buffer, whereas a blue color is emitted when tissue is processed with acidic (e.g. pH 3.3) pH buffer. It can be photographed digitally or with film (use Ektachrome 200-400 ASA film for color prints and comparable speed film for black and white prints, for example Tri-X). Most exposure times range from 10-60 second exposures, depending on the objective magnification and the intensity of the label. Thirty (30) second exposures are about average. Multiple exposures may be exploited to simultaneously visualize Fluoro-Gold and another tracer. Thus, UV would be combined with bright field illumination to simultaneously locate Fluoro-Gold with HRP or silver grains in autoradiography. Similarly, blue light excitation can be combined to also visualize the green emission color of FITC, while green excitation light may be used to simultaneously observe the red emission color of propidium iodide, or ethidium bromide (a fluorescent counterstain).

Additional Information

For pressure injections through a microsyringe or micropipette, Fluoro-Gold should be dissolved in distilled water or .9% saline. Fluoro-Gold may also be utilized as a suspension in .2M neutral phosphate buffer, however,

Contact your local distributor
Info@fluoprobes.com
Technical-support@fluoprobes.com
Order-online@fluoprobes.com

Fluoprobes, powered by



P.2

FT-T8135

the suspended particles may clog a fine micropipette tip so distilled water or .9% saline is the preferred vehicle. For iontophoresis, a 1% Fluoro-Gold solution is made up in .1M acetate buffer (pH=3.3). Well-cleaned (95% ETOH, water) glass micropipettes should have tips of 10-20 μm . Optimal iontophoresis parameters are +1 to +5 μA amps delivered with pulsed current (4-10 seconds on, 4-10 seconds off) over a 10-20 minute period.

Virtually any central or peripheral nervous system structure can be injected with Fluoro-Gold for analysis of retrograde transport. In the peripheral nervous system, ganglia and peripheral targets can be studied. For studies of peripheral nerve, the nerve should be cut or damaged and either dipped in, or injected with, aq 5% solution of Fluoro-Gold. Since Fluoro-Gold is not significantly taken up by intact fibers of passage, the fibers must be cut or severely damaged for uptake of the dye to occur. It is not presently known if Fluoro-Gold can be used for intracellular injections or used to study gap junctions, however, some groups of investigators are presently testing these possibilities.

Fluoro-Gold is used as a retrograde axonal tracer, although orthograde axonal transport does occur. The survival time should be varied (especially to very short survival times of 12 hours - 2 days) to maximize orthograde transport in the specific neuronal system under study. For retrograde transport, the survival times should be varied from 4 days to 14 days. Seven to 10 days works for most systems, although long pathways (e.g., spinal cord to brainstem) and pathways in large mammals (e.g., cats, monkeys) may require longer survival times (e.g., 14 days). In addition, since Fluoro-Gold remains fast within retrogradely labeled neurons, survival times of several months will also produce excellent results. For iontophoresis, 1 2-5 day survival time is recommended.

Tissue processing is covered in detail in the use guide and in the original publication (Schmued and Fallon, 1986, Brain Research 377:147-154). Since Fluoro-Gold is stable in many solvents and remains fast within retrogradely labeled neurons, its use is compatible with many histochemical techniques. It can be used with other retrograde tracers, immunofluorescence, PAP and ABC immunocytochemistry, HRP histochemistry, autoradiography, counterstains (ethidium bromide is the preferred fluorescent counterstain), paraffin embedding and plastic embedding. However, if tissue is unfixed, additional processing of tissue in aqueous solutions for over an hour or two will result in loss of Fluoro-Gold fluorescence from labeled neurons. Fluoro-Gold may be useful in electron microscopy. Fluoro-Gold can be used in a brain which has been sectioned and transferred to phosphate buffer. Sections are typically mounted on gelatin-coated slides, air dried, immersed in xylene and coverslipped with DPX plastic mounting media (FLUKA Chemical Corp., 255 Oser Avenue, Hauppauge, New York, 11788, Catalog #44581). Tissue may also be viewed on slides without further processing, can be run through graded alcohols for dehydration, or, for immunocytochemistry, the sections can be air dried and directly coverslipped with neutral buffered glycerine (1:2).

Fluoro-Gold is visualized with a fluorescence microscope using a wide band ultraviolet (UV) excitation filter. Use the same filter pack you would for other fluorescent retrograde tracers excited under wide band UV (e.g., True Blue, Fast Blue, Nuclear Yellow), such as the Leitz Ploem filter system A (Wide Band UV, Excitation filter BP 340-380), Mirror RKP 400, Barrier Filter LP 430). Objectives should be made especially for fluorescence microscopy (such as that made by Zeiss) glycerine, or water. Since plastic does absorb UV light, it is not advised to view through plastic petri dishes, etc. Recommended films are T-Max (Kodak, black & white) and Ektachrome 200 (Kodak, color slides). Exposure times usually vary from 20 seconds to 1.5 minutes. It is noteworthy that a photobleaching of Fluorogold reported initially due to the immunohistochemical processing, is in fact due to the water in the sections. This problem is easily corrected by using a 29:1 or 19:1 glycerine-PBS buffer mounting medium.

Fluorogold is provided as golden-yellow crystalline powder, with highest purity. Lower purity compounds, and other salts (methanethiosulfonate) have been shown to have very different properties, as solubility a bioactivity). It can be solubilized easily at 5% w/v in aqueous solutions. A 1% solution has a pH between 4.0 and 5.2 at 25°C.

The spectral characteristics vary with pH. It has a wide emission bandwidth in water, $\lambda_{\text{abs.}}/\lambda_{\text{em.}}$ [2%, pH4.5] = 323/408nm; EC: 33 000 $\text{M}^{-1} \text{cm}^{-1}$ with a second peak at ~600 nm. Spectra differ greatly when bound to membranes at a physiological pH of 7.4 ($\lambda_{\text{exc}}\lambda_{\text{em}} = 350\text{-}395 / 530\text{-}600 \text{ nm}$), to DNA ($\lambda_{\text{exc}}\lambda_{\text{em}} = 360 / \sim 450, \sim 625 \text{ nm}$; EC: 27 000 $\text{M}^{-1} \text{cm}^{-1}$), or to RNA.

FluoroGold exhibits AT-selective binding/II structure. Complex fluorescent properties allow DNA/RNA distinction.

FT-T8135

References

- **Puigdellivol-Sanchez A, et al.** « On the use of fast blue, fluoro-gold and diamidino yellow for retrograde tracing after peripheral nerve injury: uptake, fading, dye interactions, and toxicity.», *Neurosci Methods* **115**, 115-27 (2002)
- **Ju G, et al.** ; *J Neurosci Methods*. 29(1):69-72 (1989)

Related products

- Histochoice Mounting media #41927A
- Ethidium bromide #FP-06022A, 5x1g
- Propidium iodide #FP-31238B, FP-36774A
- Anti FluoroGold, #586235
- [NeuroVue](#) for Neuronal Tract Tracing

Ordering information

Catalog size quantities and prices may be found at

http://www.interchim.com/interchim/inter_intro_uptima_catalog.htm.

Please inquire for higher quantities (availability, shipment conditions).

For any information, please ask : Uptima / Interchim; Hotline : +33(0)4 70 03 73 06

Disclaimer : Materials from Uptima are sold **for research use only**, and are not intended for food, drug, household, or cosmetic use. Fluoprobes® is not liable for any damage resulting from handling or contact with this product.

FluoroGold® is a Trademark of Fluorochrome

Rev.G01E