



AZA530

Monoclonal Antibody to CD11b (Human) Phycoerythrin (PE) conjugated

CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.

Cat# : AZA530 100 tests (2ml)

Clone: MEM-174

Isotype: Mouse IgG2a

Specificity: The antibody MEM-174 recognizes CD11b antigen (Mac-1 alpha), a 165-170 kDa type I transmembrane protein mainly expressed on monocytes, granulocytes and NK-cells. The CD11b mediates neutrophil and monocyte interactions with stimulated endothelium.

Immunogen: Human granulocytes

Application: The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.
The content of a vial (2 ml) is sufficient for 100 tests.

Preparation: The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.
Do not use after expiration date stamped on vial label.
Short-term exposure to room temperature should not affect the quality of the reagent.

References:

- *Hentzen ER, Neelamegham S, Kansas GS, Benanti JA, McIntire LV, Smith CW, Simon SI. Sequential binding of CD11a/CD18 and CD11b/CD18 defines neutrophil capture and stable adhesion to intercellular adhesion molecule-1. *Blood*. 2000 Feb 1;95(3):911-20.
- *Lawrence PK, Srikumaran S.: CD11b of *Ovis canadensis* and *Ovis aries*: molecular cloning and characterization. *Vet Immunol Immunopathol*. 2007 Oct 15;119(3-4):287-98.
- *Akramiene D, Kondrotas A, Didziapetriene J, Kevelaitis E.: Effects of beta-glucans on the immune system. *Medicina (Kaunas)*. 2007;43(8):597-606.
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- *Drbal K, Moertelmaier M, Holzhauser C, Muhammad A, Fuertbauer E, Howorka S, Hinterberger M, Stockinger H, Schütz GJ: Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement. *Int Immunol*. 2007 May;19(5):675-84.

For in vitro research use only

LIFE SCIENCES

**interchim**211 bis, avenue JF Kennedy
BP 1140 - 03100 Montluçon
Fax +33 4 70 03 82 60

Hotline +33 4 70 03 73 06 • interbiotech@interchim.com