Product Name

Specific IgG against Bisphenol A Rabbit K66, titer 15,000 (at 50 % binding)

Description

Rabbit Immunoglobulin G (Caprylic acid purified) against Bisphenol A stored in 0.1 M Phosphate Buffered Saline (PBS) pH 7.2. (0.02% sodium azide)

Immunogen

BSA-Bisphenol Valeric Acid (BVA)

Cross Reactivity in a direct assay

Molecules containing	
a phenolic group	% Cross Reactivity
Bisphenol A	100%
4,4'-(ethylidene) bisphenol	10%
Bis-(4-hydroxy phenyl)-methane	1%
Nonylphenol	1%
4-cumylphenol	20%
$\Lambda I = 1 = 1 = 1 = 1 = 1 = 1 = 1 = 1 = 1 =$	
Molecules lacking	
a phenolic group	% Cross Reactivity
0	% Cross Reactivity 0.1%
a phenolic group	0
a phenolic group Vinclozolin	0.1%
<i>a phenolic group</i> Vinclozolin Pirimifos-ethyl	0.1% < 0.1%
<i>a phenolic group</i> Vinclozolin Pirimifos-ethyl 17ß-Estradiol	0.1% < 0.1% < 0.1%

Cross Reactivity in an indirect assay

J	5
Molecules containing	
a phenolic group	% Cross Reactivity
Bisphenol A	100%
4,4 ['] -(ethylidene) bisphenol	100%
Bis-(4-hydroxy phenyl)-methane	1%
4-cumylphenol	100%

Molecules lacking	
a phenolic group	% Cross Reactivity
Vinclozolin	<0.1%
Pirimifos-ethyl	< 0.1%
2,4 D	0.1%
Fenitrothion	< 0.1%
Chlorpyrifos-methyl	< 0.1%
Erythromycine	< 0.1%

Specificity

This antibody is highly specific for Bisphenol A

Tested applications

ELISA

Application notes Suggested concentration to use in ELISA : 1/15,000 from the delivered solution. HRP-conjugated derivative as a tracer

Relevance

Bisphenol A is a known hormone-disrupting agent, commonly used in plastics and diffusing into the environment and food.

Raised in Rabbit

Clonality Polyclonal

Storage buffer Phosphate buffered saline

Concentration

Appr. 0.8 mg/ml

Storage instructions

This working antibody solution is stable for at least 7 days at 4 °C. Precautions of storage should be taken for longer periods. Problems of long term stability may occur with highly diluted solutions. No other preservative agent has been added to the present formulation. For long storage purposes in solution the addition of sodium azide at 0.02 % is advised with the appropriate precautions of use.

References

Eline P. Meulenberg, Kees Koopal and Ria Rhemrev - Immunoassays for alkylphenolic pollutants with endocrine disrupting activity. Intern. J. Environ. Anal. Chem. Vol 85, no 12-13, 15 October-15 November 2005, 871-883