

TEMPase Hot Start DNA Polymerase Glycerol Free

Concentration: 5 units/ μ l



Cat. No.: A240003

500 Units

	TEMPase Polymerase, Glycerol Free, 5 U/ μ l
ID No.	5101700
Cap colour	White
A240003	100 μ l

MADE IN DENMARK

Key Features

TEMPase Hot Start DNA Polymerase, Glycerol Free, is a high-quality DNA polymerase developed for automation. It is a glycerol free formulation of TEMPase Hot Start Polymerase with the same excellent performance. The glycerol free formulation makes it well suited for automation and freeze-drying.

TEMPase Hot Start DNA Polymerase is a modified form of Ampliqon Taq DNA Polymerase. A chemical moiety is attached to the enzyme at the active site, which renders the enzyme inactive at room temperature. Thus, during setup and the first ramp of thermal cycling, the enzyme is not active and misprimed primers are not extended. Once the reaction reaches optimal activating temperature, the chemical moiety is cleaved during a 15 minutes heat activation step, releasing the active TEMPase Hot Start DNA Polymerase into the reaction.

Kit Components

TEMPase Hot Start DNA Polymerase in Glycerol Free Storage Buffer

5 U/ μ l TEMPase Hot Start DNA Polymerase, 20 mM Tris-HCl pH 8.9, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween[®] 20.

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

TEMPase Hot Start DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Unit Definition

One unit is defined as the amount of polymerase that incorporates 10 nmol of dNTPs into acid-precipitable DNA in 30 minutes at 72 °C under standard assay conditions.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using TEMPase Hot Start DNA Polymerase, Glycerol free. Optimal reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually.

1. Thaw Solutions. **It is important to thaw all solutions completely (some buffers need to reach room temperature) and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice.
2. Prepare a master mix according to Table 1. The master mix typically contains all the components needed for extension except the template DNA.

Table 1. Reaction mix and template DNA

Component	Vol./reaction*	Final concentration*
10x Buffer	5 μ l	1x
25 mM MgCl ₂	-	Optional
dNTP mix (10 mM each)	1 μ l	0.2 mM of each dNTP
Primer A (10 μ M)	1 μ l (0.5 – 5 μ l)	0.2 μ M (0.1 – 1.0 μ M)
Primer B (10 μ M)	1 μ l (0.5 – 5 μ l)	0.2 μ M (0.1 – 1.0 μ M)
TEMPase DNA Pol.	0.4 μ l (0.2 – 1 μ l)	2 units (1 – 5 units)
PCR-grade H ₂ O	X μ l	-
Template DNA	X μ l	genomic DNA: 20 ng (1 – 200 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng)
TOTAL volume	50 μ l	-

* Suggested starting conditions; theoretically used conditions in brackets. The final volume can be reduced to 25 μ l by using half of the volumes suggested in Vol./reaction, eg. 0.2 μ l TEMPase instead of 0.4 μ l TEMPase.

3. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes.
4. Add template DNA to the individual tubes containing the reaction mix.
5. Program the thermal cycler according to the manufacturer's instructions. **Each program must start with an initial heat activation step at 95°C for 15 minutes.** For maximum yield and specificity, temperatures and cycling times should be optimized for each new template or primer pair.
6. Place the tubes in the thermal cycler and start the reaction.

Table 2. Three-step PCR program

Cycles	Duration of cycle	Temperature
1	15 minutes ^a	95 °C
25 – 35	20 – 30 seconds ^b 20 – 40 seconds ^c 30 – 90 seconds ^d	95 °C 50 – 65 °C 72 °C
1	5 minutes ^e	72 °C

^a For activation of the TEMPase hot start enzyme.

^b Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^c Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-

stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used.

- ^d Extension/elongation step: TEMPase polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.
- ^e Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Tween® 20 is a registered trademark of ICI Americas Inc.

Notes:

- For longer DNA targets more DNA polymerase could be added to the PCR master mix.

Related Products

TEMPase DNA Polymerase (500 units) *	Cat. No.
TEMPase Hot Start DNA Polymerase 5 U/μl	A220003
• with 10x Ammonium Buffer	A221103
• with 5x PCR Buffer RED	A221803
TEMPase Hot Start DNA Polymerase 5 U/μl, glycerol free	A240003
• with 10x Ammonium Buffer	A241103

Taq DNA Polymerase (500 units) *	Cat. No.
Taq DNA Polymerase, 5 U/μl	A110003
• with 10x Ammonium Buffer	A111103

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM MgCl₂.

Buffers for DNA polymerases *	Cat. No.
10x Ammonium Buffer, 3 x 1.5 ml	A301103
10x Standard Buffer, 3 x 1.5 ml	A302103
10x Combination Buffer, 3 x 1.5 ml	A303103
5x PCR Buffer RED, 6 x 1,5 ml **	A301810

*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as Mg²⁺ free buffers, detergent free buffers and Mg²⁺ and detergent free buffers.

**For direct gel loading and visualisation.

TEMPase Hot Start Master Mixes (500 x 50 μl reactions) *	Cat. No.
2x Master Mix A**, 1.5 mM MgCl ₂ final concentration	A230303
2x Master Mix A**BLUE, 1.5 mM MgCl ₂ final concentration	A290403

*Master mixes available also in 1.1x variants as well as 2 mM MgCl₂ variants, **Mix A is Ammonium Buffer based, also available as Mix C based on Combination Buffer.

Special Master Mixes (500 x 50 μl reactions)	Cat. No.
Multiplex 2x Master Mix, 3 mM MgCl ₂ final concentration	A260303
GC TEMPase 2x Master Mix I – for GC-rich templates	A331703
GC TEMPase 2x Master Mix II – for GC-rich templates	A332703

Ultrapure dNTPs*	Cat. No.
dNTP Mix 40 mM (2 x 500 μl): 10 mM each dA, dC, dG, dT	A502004
dNTP Set, 100 mM each: 250 μl of each dA, dC, dG and dT	A511104

*Other concentrations and Single dNTPs are available.

Loading Buffers, PCR water and Ladders	Cat. No.
5x Loading Buffer Red *, 5 x 1 ml	A608104
Iqon PCR Ladder **, 100 – 3000 bp, 1 x 0.5 ml	A610341
PCR Grade Water, 6 x 5 ml	A360056

* Also available with Blue, Orange or Cyan. ** Available in different size ranges.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 08/2021

