

Latex Conjugation Kit – Mini Vial Conjugation

Applicable to: **Blue Latex:** 1000-0040, 1000-0100
Red Latex: 1002-0040, 1002-0100
Black Latex: 1004-0040, 1004-0100

Release 2. © EXPEDEON. 23/11/2018


INTRODUCTION

The Latex Conjugation Kit allows antibodies or proteins to be conjugated to our high quality 400 nm latex nanoparticles quickly and easily. The latex nanoparticles are designed for ease of use and have much improved handling compared with traditional latex. This has been achieved by specially treating the nanoparticles. The conjugation is covalent, and requires less antibody than traditional passive and covalent latex conjugation.

The 400 nm blue, red or black latex nanoparticles in this kit are freeze dried. The conjugation reaction is initiated simply by reconstituting the dry mixture with your antibody, which becomes attached (via lysine residues) to the specially treated surface.

It takes 30 seconds to set up the conjugation, the hands-on time for the conjugation procedure is about 3 minutes and the conjugate is ready to use within 35 minutes. You simply pipette the biomolecule into a vial containing the latex nanoparticles, then centrifuge to buffer exchange (Figure 1).

Figure 1. Latex Conjugation



The resulting covalent conjugates can be easily resuspended without the need for harsh methods such as sonication or vortexing, unlike traditional conjugation procedures which are prone to aggregation. This is due to the properties of the surface treatment which makes the particles resistant to aggregation.

Additionally, unlike passive methods, the conjugation procedure has only a weak dependence on the isoelectric point of the antibody. Consequently, extensive trials at different pH values are not required; all antibodies can be conjugated at one of two pHs, both of which are supplied in this kit (Reaction Buffers A and B).

We would suggest that the antibody to be conjugated should be in 10 – 50 mM MES, HEPES or MOPS at pH 6 – 7 (with no other components e.g. salt or azide) before conjugation. See “Buffer Considerations” for more details and advice if your antibody is in another buffer.

KIT CONTENTS

- 4 or 10 Mini vials of 400 nm Blue, Red or Black latex (dependent on kit)
Each Mini vial gives sufficient conjugate for ~50 lateral flow tests
- 1 vial of 1x Reaction Buffer A
- 1 vial of 1x Reaction Buffer B
- 1 vial of 10x Quencher
- 1 vial of Resuspension Buffer

SHIPPING CONDITIONS

The kit is shipped at ambient temperature in a tamper-evident polypropylene container. Please store the kit at -20°C upon receipt. All the buffers and the Quencher can be stored at either 4°C or -20°C.

INSTRUCTIONS

1. Allow all of the reagents to warm to room temperature
2. Dilute your stock antibody to 0.1mg/ml with Reaction Buffer A or B. 45µl will be sufficient to conjugate one Mini vial.
NOTE: *If this is the first test of an antibody we advise carrying out the reaction twice, once with Reaction Buffer A and once with Reaction Buffer B to find the optimal pH. If you also wish to examine the effect of varying the amount of antibody, make different antibody dilutions.*
3. Add 40µl of the 0.1mg/ml antibody to the Mini vial and reconstitute the latex nanoparticles by gently and thoroughly pipetting up and down. Incubate the reaction for exactly 15 minutes at room temperature.
4. Dilute sufficient 10x Quencher with deionised water. For one Mini vial you need exactly 1ml so we advise you to make 1.2ml 1x Quencher per vial i.e. 120µl 10x Quencher + 1080µl water.
5. After 15 minutes, add 1ml 1x Quencher to stop the reaction, and mix by inverting several times.
6. Leave the reaction to quench for 5 minutes at room temperature, then transfer into a microcentrifuge tube and spin at 10,000 rpm for 9 minutes. Remove ~850µl of the supernatant and without resuspending the pellet spin for 1 more minute at 10,000 rpm. Remove the rest of the supernatant.
7. Gently resuspend the pellet in 40µl Resuspension Buffer or a storage buffer of your choice.
NOTE: *For increased stability of the liquid conjugate we advise adding 0.1% BSA (final concentration) to the Resuspension Buffer just prior to use. e.g. add 2.5µl 2% BSA diluted in deionised water to 50 µl Resuspension Buffer.*
8. You now have 40µl 1% conjugate. Dilute further as required for your application.

AMOUNT AND VOLUME OF ANTIBODY

The optimum amount of antibody (which will influence the number of antibody molecules per particle) depends on the size of the nanoparticles (surface area) and on the application; you may need to conjugate different amounts of antibody to optimize your assay.

We recommend testing with 0.1 mg/ml in the first instance, although slightly lower or higher concentrations can be explored to optimize performance in your application.

Before testing antibody concentrations above 0.1 mg/ml we advise you to use our Antibody Concentration & Clean Up Kit for Latex and Europium to remove interfering buffer components (see buffer considerations below). This is due to the increase in contamination that will occur when conjugating using a larger volume of the stock antibody.

Do not alter the reaction volume of 40µl as this will reduce conjugation efficiency.

STORAGE OF CONJUGATES

Initial conjugate storage at 4°C is recommended.

The Resuspension Buffer added at the end of the conjugation reaction is a good storage buffer. **Do not store the conjugate at -20°C.**

The determining factor for conjugate stability will be the antibody itself, as it will be first to degrade. Therefore, as long as your antibody is stable, the conjugate will be stable as well.

BUFFER CONSIDERATIONS

There are a number of common buffer components that have a substantial negative effect on the conjugation efficiency. This decreases the amount of antibody that will be coupled to each nanoparticle, so reduces the signal from and sensitivity of the conjugated latex. To prevent this we advise conjugating only from stock antibodies that are at least 1 mg/ml in 10 – 50 mM MES, HEPES or MOPS at pH 6 – 7 (with no other components e.g. salt or azide).

These are not common antibody storage buffers, so we have developed the Antibody Concentration & Clean Up Kit for Latex and Europium for use with antibodies stored in other buffers and therefore not compatible with this kit. This kit will quickly and simply purify your antibody into Reaction Buffer A and/or B. Please see the kit protocol for more details.

Please see the table for a list of compatible and incompatible buffer components for this kit, and those that can be removed by the Antibody Concentration & Clean Up Kit for Latex and Europium. Proteins cannot be removed by the Antibody Concentration & Clean Up Kit for Latex and Europium, however our AbPure™ kits can be used to achieve this.

<i>Buffer components</i>	<i>Latex and Europium Conjugation Kit can tolerate</i>	<i>Concentration & Clean Up Kit can remove</i>
pH 6 - 7	✓	✓
pH < 6 and > 7	✗	✓
Amine free buffer (≤50 mM) (e.g. MES, MOPS, HEPES)	✓	✓
Amine free buffer (≥50 mM) (e.g. MES, MOPS, HEPES)	✗	✓
Salt	✗	✓
Sodium Azide	✗	✓
Sugars	✓	✓
Glycerol	✓	✓
Thiomersal	✗	✓
Thimerosal	✗	✓
Merthiolate	✗	✓
BSA	✗	✗ ¹
Gelatin	✗	✗ ¹
Tris	✗	✓
Glycine	✗	✓
Carboxylic acids (e.g. EDTA, Citrate)	✗	✓
Nucleophilic components (Primary amines e.g. amino acids or ethanolamine and thiols e.g. mercaptoethanol or DTT)	✗	✓

¹ If the antibody to be conjugated contains other proteins such as BSA or Gelatin we recommend using our range of AbPure™ kits to remove such unwanted proteins

TECHNICAL SUPPORT

For technical enquiries get in touch with our technical support team at: www.expedeon.com/contact/

Limited use license

Use and Use Restrictions: The Products are sold, and deliverables of any services are provided for the purposes of the buyer's internal in vitro research, development or educational use only, not for in vivo, or any therapeutic or diagnostic use, nor for resale, or for providing services or any other commercial use of any kind, including without limitation, for any transfer in any form (including as part of a kit) to a third party; for analysis or reverse engineering of the Product or for manufacturing. Products should only be used in accordance with any safety data sheets, guidance or protocols that we issue from time to time and are available for download from our Site. Protective clothing should be used at all times when handling our Products. Safety datasheets relating to all Products are available for download from the Site or upon request.

Expedeon grants no other license or rights under any intellectual property in respect of Products or services deliverables and in particular grants no license to use any Product or deliverables for any commercial purposes.

Sale of Products or service deliverables by us or our authorised distributors are expressly conditional upon the customer's agreement with these restrictions, which the customer gives upon placing an

order for Products or deliverables. If you wish to use any Product or deliverables for any purpose other than your own internal research as described above, you will require an additional licence from Expedeon. Please contact licensing@expedeon.com