

FT-80910



Brain Heart Infusion Agar (BHI Agar)

Recommended for the development of fastidious microorganisms

Product Description

Name :	Brain Heart Infusion Agar (BHI Agar)			
Catalog Number :	809108, 500 g			
Formula (g/l) :	Peptone Mixture	10.00	Disodium Phosphate	2.50
	Beef Heart Infusion	10.00	Sodium Chloride	5.00
	Calf Brain Infusion	7.50	Bacteriological Agar	15.00
	Dextrose	2.00		
pH:	Final pH 7.4 ± 0.2 at 25°C			

Storage: +2°C to 25°C
Protect from light and moisture. Once opened keep powdered medium closed to avoid hydration.

Directions for use

Preparation

Suspend 52 grams of the medium in one liter of distilled water. Mix well and dissolve by heating with frequent agitation. Boil for one minute until complete dissolution. Sterilize in autoclave at 121°C for 15 minutes. The prepared medium should be stored at 8-15°C. The color is clear amber, slightly opalescent.

The dehydrated medium should be homogeneous, free-flowing and beige in color. If there are any physical changes, discard the medium.

To prepare a selective medium for fungi, the sterilized and melted medium should be cooled to 45 - 50°C before adding the appropriate antibiotics.

Occasionally a small amount of sediment may appear which should be resuspended with a gentle swirl before dispensing.

Uses

BRAIN HEART INFUSION AGAR (BHIA) is a solid medium rich in nutrients, suitable for the cultivation of several fastidious strains of bacteria, fungi, and yeasts.

Brain Heart Infusion Agar is used for the cultivation of a wide variety of fastidious microorganisms such as streptococci, meningococci and pneumococci. BHIA is recommended in Standard Methods for water testing and in antimicrobial susceptibility tests. The nutritionally rich base of Beef heart and Calf brain infusions and Peptone mixture provide nitrogen, vitamins, minerals and amino acids that supports the growth of a variety of microorganisms.



FT-80910

Disodium phosphate acts as a buffer. Dextrose is the fermentable carbohydrate providing carbon and energy. Sodium chloride maintains the osmotic balance. Bacteriological agar is the solidifying agent.

Inoculate and incubate anaerobically at $35 \pm 2^\circ\text{C}$ for 24 - 72 hours. If 10% sterile defibrinated blood is added, the medium can be used for the cultivation and isolation of *Histoplasma capsulatum*. With the addition of antibiotics the medium can be used for the isolation of fungi. Brain Heart Infusion Agar with cycloheximide and chloramphenicol restrict growth of bacteria and saprophytic fungi, and is recommended for the isolation of fungi difficult to grow such as *H. capsulatum* and *Blastomyces dermatitidis*. Adding polysorbate to BHIA allows for identification of *Mycobacterium avium* - intracellulare complex organisms and *M. tuberculosis* from blood cultures.

Occasionally BHIA plates are used for general sensitivity tests. However, it is not suitable to determine hemolytic reactions as this medium has a high dextrose concentration and it may give atypical readings.

Microbiological test

The following results were obtained in the performance of the medium from type cultures after incubation at a temperature of $35 \pm 2^\circ\text{C}$, under 5 - 10% CO_2 , and observed after 24 - 72 hours. (It is recommended to grow *Aspergillus brasiliensis* and *Saccharomyces cerevisiae* aerobically at $30 \pm 2^\circ\text{C}$).

Microorganisms	Growth without blood	Growth with 5% sheep blood
<i>Aspergillus brasiliensis</i> ATCC 16404	Good	Good
<i>Neisseria meningitidis</i> ATCC 13090	Moderate	Good
<i>Saccharomyces cerevisiae</i> ATCC 9763	Good	Good
<i>Streptococcus pneumoniae</i> ATCC 6303	Moderate	Good
<i>Streptococcus pyogenes</i> ATCC 19615	Moderate	Good

Bibliography

- Creitz and Pucket A.J. Clin. Path 24:1318, 1954.
- Golding and Davidson Modern, Hospital, 92:April 1954

Ordering information

Catalog size quantities and prices may be found at <http://www.interchim.com>. Please inquire for higher quantities (availability, shipment conditions).

For any information, please ask : Uptima / Interchim; Hotline : +33(0)4 70 03 73 06
[Order on-line](#) or [Contact](#) your local distributor

Disclaimer : Materials from Uptima are sold **for research use only**, and are not intended for food, drug, household, or cosmetic uses. Uptima is not liable for any damage resulting from handling or contact with this product.