

FT- JQ7610



INTERFINE CHEMICALS ANALYTICAL SCIENCES BIOCHROMATOGRAPHY BIOSCIENCES


Annexin V, recombinant

To measure swiftly apoptosis in a variety of suspended cell types.

Product Information

Name :	AnnexinV , recombinant
Catalog Number :	FP-JQ7610, 100µg FP-JQ7611, 1mg
Concentration :	2 mg/ml in PBS (pH 7,4)
Solubility :	Water
MW :	~36 kD
Source :	<i>E. coli</i>
Purity :	>95% by SDS-PAGE
Storage:	-80°C

Expiration date is 6 months from the date of receipt.



Technical information

Annexin V is used as a probe in annexin V affinity assay to detect apoptotic cells, which have expressed phosphatidylserine on the cell surface, a feature found in apoptosis as well as other forms of cell death. Apoptosis is a form of programmed cell death, which is used by the body to remove unwanted, damaged or senescent cells from tissues. Removal of apoptotic cells is carried out via phagocytosis by white blood cells such as macrophages or dendritic cells.

Annexin V has been successively used to detect apoptotic cells *in vitro* (cells in a culture tube) and *in vivo* (in laboratory mice and in patients). Pathological processes in which apoptosis occurs include inflammation, ischemia damage of the heart cause by myocardial infarction, apoptotic white blood cells present in an atherosclerotic plaque in blood vessels, transplanted organs in the donor patient which are rejected by the immune system or tumor cells which are exposed to cytostatic drugs during chemotherapy.

Phagocytic white blood cells recognize apoptotic cells by their exposure of negatively charged phospholipids (phosphatidylserine) on the cell surface. In normal cells the negative phospholipids reside on the inner side of the cellular membrane and the outer surface of the membrane is occupied by phospholipids, which do not have a charge. After cells have made the decision to commit suicide, the negatively charged phospholipids are transported to the outer cell surface by a hypothetical protein known as scramblase. Phagocytic white blood cells express a receptor, which can detect the negatively charged phospholipids. After detection the apoptotic cells are removed. Phagocytes rapidly remove the apoptotic cells of healthy individuals. However in pathological processes the removal of apoptotic cells may be delayed or even absent. Dying cells in tissue can be detected with annexin V.

FT- JQ7610

Labeling of annexin V with fluorescent or radioactive molecules makes it possible to detect binding of labeled annexin V to the cell surface of apoptotic cells. After binding to the phospholipid surface, annexin V assembles into a cluster known as a trimer. This trimer consists of three annexin V molecules, which are bound to each other via non-covalent protein-protein interactions. The formation of annexin V trimers results in the formation of a two-dimensional crystal lattice on the phospholipid membrane. This clustering of annexin V on the membrane greatly increases the intensity of annexin V when labeled with a fluorescent or radioactive probe.

Annexin V is available as conjugated to FluoProbes, biotin, FITC and more. See related products or inquire at info@fluoprobes.com.

More information in technical sheet of AnnexinV-FluoProbes® 488 [FP- BH4140](#) (apoptosis, directions for use, references).

Related products

- 7-Aminoactinomycin (7-AAD), [FP-132303](#)
- Propidium iodide, 1 mg/ml, [FP-36774A](#)
- AnnexinV binding buffer, [FP-BU2080](#)
- Annexin V-FluoProbes® 488, *in vivo* Grade, [FP-BY6820](#)
- Annexin V-FluoProbes® 488, *Confocal microscopy* Grade, [FP- BH4140](#)

Ordering information

Please inquire for higher quantities (availability, shipment conditions).

For any information, please ask : FluoProbes® / Interchim; Hotline : +33(0)4 70 03 73 06

Disclaimer : Materials from FluoProbes® are sold **for research use only**, and are not intended for food, drug, household, or cosmetic use. FluoProbes® is not liable for any damage resulting from handling or contact with this product.

Rev.I11E